

References for articles indicating the benefits of physiotherapy

Below, please find a list of references to articles which indicate the benefits of physiotherapy. You can use these to back up articles and publicity about the profession, and what it achieves, particularly in the field of preventing and treating diseases of civilisation.

Hypertension

ACSM's Guidelines for Exercise Testing and Prescription. 6th Ed. Baltimore MD: Lippincott Williams & Wilkins 2000 <http://www.aafp.org/afp/20050601/2123.html>

Blumenthal JA, Sherwood A, et al. Exercise and weight loss reduce blood pressure in men and women with mild hypertension: effects on cardiovascular, metabolic, and hemodynamic functioning. Arch Intern Med. 2000;160(13): 1947-58.

Miller ER, Erlinger TP, Young DR, et al. Results of the diet, exercise, and weight loss intervention trial. Hypertension. 2002;40(5):612-618.

Tanaka H, Bassett DR, Howley ET, Thompson DL, Ashraf M, Rawson FL. Swimming training lowers the resting blood pressure in individuals with hypertension. J Hypertens. 1997;15:651-7.

Cardiovascular disease

Ades P. Cardiac Rehabilitation and Secondary Prevention of Coronary Heart Disease. N Eng J Med. 2001; 345, 12.

Balady G et al. Cardiac rehabilitation programs. A statement for healthcare professionals from the American Heart Association. Circ. 1994;90:1602-10.

Brubaker PH, Kaminsk LK, Whaley MH. Coronary Artery Disease: Essentials of Prevention and Rehabilitation Programs. Champaign Il: Human Kinetics, 2002.

Brubabaker PH, Warner JG, Rejeski DG, et al. Comparison of standard and extended length participation in cardiac rehabilitation on body composition, functional capacity, and blood lipids. Am J Cardiol 1996;78:769-773.

Frownfelter D, Dean E. Cardiovascular and Pulmonary Physical Therapy. 4th Edition. St. Louis. Mosby Elsevier. 2006.

WCPT Secretariat

Kensington Charity Centre • 4th Floor Charles House • 375 Kensington High Street • London W14 8QH • UK
T +44 (0)20 7471 6765 • F +44 (0)20 7471 6766 • info@wcpt.org • www.wcpt.org

Pollock M et al. Resistance Exercise in Individuals With and Without Cardiovascular Disease: An Advisory From the Committee on Exercise, Rehabilitation, and Prevention, Council on Clinical Cardiology, American Heart Association. *Circ*. 2000; 101: 828.

Seki E et al. Effects of Phase III Cardiac Rehabilitation Programs on Health-related Quality of Life in Elderly Patients with Coronary Artery Disease. *Circ J*. 2003; 67: 73-77.

Vincent K, Vincent H. Resistance Training for Individuals With Cardiovascular Disease. *J Cardiopulm Rehab*. 2006; 26: 207-16.

Stroke

Dean CM, Richards CL, Malouin F. Task-related circuit training improves performance of locomotor tasks in chronic stroke: a randomized, controlled pilot trial. *Arch Phys Med Rehabil*. 2000;81(4):409-17.

Endres M, Gertz K, et al. Mechanisms of stroke protection by physical activity. *Ann Neurol*. 2003;54(5):582-90.

Ouellette MM, LeBrasseur NK, et al. High-intensity resistance training improves muscle strength, self-reported function, and disability in long-term stroke survivors. *Stroke*. 2004;35(6):1404-9.

Teixeira-Salmela LF, Olney SJ, et al. Muscle strengthening and physical conditioning to reduce impairment and disability in chronic stroke survivors. *Arch Phys Med Rehabil*. 1999;80(10):1211-8.

Diabetes

Castaneda C, Layne JE, Munoz-Orians L, et al. A randomized controlled trial of resistance exercise training to improve glycemic control in older adults with type 2 diabetes. *Diabetes Care*. 2002;25(12):2335-41.

Dunstan DW, Daly RM, Owen N, Jolley D, De Courten M, Shaw J, Zimmet P. High-intensity resistance training improves glycemic control in older patients with type 2 diabetes. *Diabetes Care*. 2002 Oct;25(10):1729-36.

Evans WJ. Effects of exercise on body composition and functional capacity of the elderly. *J Gerontol A Biol Sci Med Sci*. 1995;50 Spec No:147-50.

Fenicchia LM, Kanaley JA, Azevedo JL Jr, et al. Influence of resistance exercise training on glucose control in women with type 2 diabetes. *Metabolism*. 2004;53:284–289.

Obesity

Dunstan DW, Daly RM, Owen N, Jolley D, De Courten M, Shaw J, Zimmet P. High-intensity resistance training improves glycemic control in older patients with type 2 diabetes. *Diabetes Care*. 2002 Oct;25(10):1729-36.

Hagberg JM, Graves JE, Limacher M, et al. Cardiovascular responses of 70- to 79-yr-old men and women to exercise training. *J Appl Physiol.* 1989;66(6):2589-94.

Chronic Obstructive Pulmonary Disease

American College of Sports Medicine. *Guidelines for Exercise Testing and Prescription.* 7th ed. Champaign IL: Human Kinetics, 2006.

Casaburi R, Patessio A, Ioli F, et al. Reduction in exercise lactic acidosis and ventilation as a result of exercise training in patients with obstructive lung disease. *Am Rev Respir Dis.* 1991;143:9-18.

Casaburi R, Porszasz J, Burns MR, Carithers ER, et al. Physiologic benefits of exercise training in rehabilitation of severe COPD patients. *Am J Respir Crit Care Med.* 1997;155:1541–1551.

Casaburi R. Mechanisms of the reduced ventilatory requirement as a result of exercise training. *Eur Respir Rev.* 1995;5:25, 42–46.

Clark CJ, Cochrane LM, et al. Skeletal muscle strength and endurance in patients with mild COPD and the effects of weight training. *Eur Respir J.* 2000;15(1):92-97.

Coppoolse R, Schols A, Baarends EM et al. Interval versus continuous training in patients with severe COPD: a randomized clinical trial. *Eur Respir J.* 1999;14:258-263.

Gosselink R, Troosters T, Decramer M. Effects of exercise training in COPD patients: interval versus endurance training. *Eur Respir J.* 1998;12:2S.

Gosselink R, Troosters T, Decramer M. Exercise training in COPD patients: the basic questions. *Eur Respir J.* 1997;10:2884–2891.

Griffiths TL, Burr ML, Campbell IA, et al. Results at 1 year of outpatient multidisciplinary pulmonary rehabilitation: a randomized controlled trial. *Lancet.* 2000;355:362-368.

Hernandez MTE, Rubio TM, Ruiz FO, et al. Results of a home-based training program for patients with COPD. *Chest.* 2000;118:106-114.

Hirata K, Okamoto T, Shiraishi S. The efficacy and practice of exercise training in patients with chronic obstructive pulmonary disease (COPD). *Nippon Rinsho.* 1999;57(9):2041-5.

Lacasse Y, Brosseau L, Milne S, et al. Pulmonary rehabilitation for chronic obstructive pulmonary disease, *Cochrane Review.* The Cochrane Library, Issue 3, 2002. Oxford: Update Software.

Maltais F, LeBlanc P, Jobin J, et al. Intensity of training and physiologic adaptation in patients with chronic obstructive pulmonary disease. *Am J Respir Crit Care Med.* 1997;155:555–561.

Normandin EA, McCusker C, Connors ML, et al. An evaluation of two approaches to exercise conditioning in pulmonary rehabilitation. *Chest.* 2002;121:1085-1091.

Puhan MA, Scharplatz M, Troosters T, et al. Respiratory rehabilitation after acute exacerbation of COPD may reduce risk for readmission and mortality: a systematic review. *Respir Res.* 2005;6:54.

Ries AL, ACCP/AACVPR Pulmonary Rehabilitation Guidelines Panel. Pulmonary rehabilitation: joint ACCP/AACVPR evidence-based guidelines. *Chest.* 1997;112:1363-1396.

Troosters TR, Casaburi R, Gosselink R, Decramer M. Pulmonary rehabilitation in chronic obstructive pulmonary disease. *Am J Respir Crit Care Med.* 2005;172(3):19-38.

Vogiatzis I, Nanas S, Roussos C. Interval training as an alternative modality to continuous exercise in patients with COPD. *Eur Respir J.* 2002;20:12-19.

ZuWallack R. The nonpharmacologic treatment of chronic obstructive pulmonary disease: advances in our understanding of pulmonary rehabilitation. *Proc Am Thorac Soc.* 2007 Oct 1;4(7):549-53. Review.

HIV/AIDS

Barnish M, Condoluci D, Farrar JT. The effect of group aerobic exercise and t'ai chi on functional outcomes and quality of life for persons living with acquired immunodeficiency syndrome. *J Altern Complement Med.* 2005;11(6):1085-92.

O'Brain K, Nixon S, Glazier RH, Tynan AM. Progressive resistive exercise interventions for adults living with HIV/AIDS. *The Cochran Library.* 2006:1.

Smith B, Neidig J, Nickel J, Mitchell G, Para M, Fass R. Aerobic exercise: effects on parameters related to fatigue, dyspnea, weight and body composition in HIV-infected adults. *AIDS.* April 2001;15(6):693-701.

Williams B, Waters D, Parker K. Evaluation and treatment of weight loss in adults with HIV disease. *Am Acad Family Physicians.* 1999;60(3).

Copyright © World Confederation for Physical Therapy 2008.